

The Mock LISA Data Challenges: from Challenge 1B to Challenge 3

Stanislav Babak¹, John G Baker², Matthew J Benacquista³,
Neil J Cornish⁴, Jeff Crowder⁵, Shane L Larson⁶, Eric Plagnol⁷,
Edward K Porter¹, Michele Vallisneri^{5,8}, Alberto Vecchio⁹ (*The Mock
LISA Data Challenge Task Force*), Keith Arnaud², Leor Barack¹⁰,
Arkadiusz Błaut¹¹, Curt Cutler^{5,8}, Stephen Fairhurst¹²,
Jonathan Gair^{1,13}, Xuefei Gong¹⁴, Ian Harry¹², Deepak Khurana¹⁵,
Andrzej Królak¹⁶, Ilya Mandel^{8,17}, Reinhard Prix¹⁸, B S Sathyaprakash¹²,
Pavlin Savov⁸, Yu Shang¹⁴, Miquel Trias¹⁹, John Veitch⁹, Yan Wang²⁰,
Linqing Wen^{1,8,21} and John T Whelan¹ (*the Challenge-1B participants*)

¹ Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), Am Mühlenberg 1, D-14476 Golm bei Potsdam, Germany

² Gravitational Astrophysics Laboratory, NASA Goddard Space Flight Center, 8800 Greenbelt Rd., Greenbelt, MD 20771, USA

³ Center for Gravitational Wave Astronomy, University of Texas at Brownsville, Brownsville, TX 78520, USA

⁴ Department of Physics, Montana State University, Bozeman, MT 59717, USA

⁵ Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA

⁶ Department of Physics, Weber State University, 2508 University Circle, Ogden, UT 84408, USA

⁷ APC, UMR 7164, University Paris 7 Denis Diderot, 10, rue Alice Domon et Leonie Duquet, 75025 Paris Cedex 13, France

⁸ Theoretical Astrophysics, California Institute of Technology, Pasadena, CA 91125, USA

⁹ School of Physics and Astronomy, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

¹⁰ School of Mathematics, University of Southampton, Southampton, SO17 1BJ, UK

¹¹ Institute of Theoretical Physics, University of Wrocław, Wrocław, Poland

¹² School of Physics and Astronomy, Cardiff University, 5, The Parade, Cardiff, CF24 3YB, UK

¹³ Institute of Astronomy, University of Cambridge, Madingley Rd., Cambridge, CB3 0HA, UK

¹⁴ Institute of Applied Mathematics, Academy of Mathematics and System Science, Chinese Academy of Sciences, 55 Zhongguancun Donglu, Beijing 100080, People's Republic of China

¹⁵ Indian Institute of Technology, Kharagpur, India

¹⁶ Institute of Mathematics, Polish Academy of Sciences, Warsaw, Poland

¹⁷ Department of Physics and Astronomy, Northwestern University, Evanston, IL, USA

¹⁸ Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), D-30167 Hannover, Germany

¹⁹ Department de Física, Universitat de les Illes Balears, Cra. Valldemossa km 7.5, E-07122 Palma de Mallorca, Spain

²⁰ Department of Astronomy, Nanjing University, 22 Hankou Road, Nanjing 210093, People's Republic of China

²¹ School of Physics, M013, University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia

E-mail: Michele.Vallisneri@jpl.nasa.gov

Received 21 April 2008, in final form 11 June 2008

Published 2 September 2008

Online at stacks.iop.org/CQG/25/184026

Abstract

The Mock LISA Data Challenges are a programme to demonstrate and encourage the development of LISA data-analysis capabilities, tools and techniques. At the time of this workshop, three rounds of challenges had been completed, and the next was about to start. In this paper we provide a critical analysis of the entries to the latest completed round, Challenge 1B. The entries confirm the consolidation of a range of data-analysis techniques for galactic and massive-black-hole binaries, and they include the first convincing examples of detection and parameter estimation of extreme-mass-ratio inspiral sources. In this paper we also introduce the next round, Challenge 3. Its data sets feature more realistic waveform models (e.g., galactic binaries may now chirp, and massive-black-hole binaries may precess due to spin interactions), as well as new source classes (bursts from cosmic strings, isotropic stochastic backgrounds) and more complicated nonsymmetric instrument noise.

PACS numbers: 04.80.Nn, 95.55.Ym

1. Introduction

The Laser Interferometer Space Antenna (LISA), an ESA–NASA mission to survey the gravitational-wave (GW) sky at frequencies between 10^{-5} and 10^{-1} Hz, will record gravitational radiation from millions of sources, most of them in our galaxy, but many populating the low-to-high-redshift universe [1]. Such a variety of signals, overlapping in both the time and frequency domains (to the point of creating *confusion noise* at some frequencies) poses a number of interesting new challenges for GW data analysis, whose solution is essential if we are to draw the greatest possible science payoff from such a bold and innovative observatory.

At the end of 2005, the LISA International Science Team (LIST) initiated a programme of Mock LISA Data Challenges (MLDCs) with the goal of understanding at the conceptual and quantitative level the peculiarities of LISA data analysis, of demonstrating LISA’s observational capabilities, and kickstarting the development of data-analysis algorithms, pipelines and infrastructural elements. An MLDC Task Force, chartered by the LIST, periodically issues *challenge* data sets containing GW signals from sources of undisclosed parameters, embedded in synthetic LISA noise; challenge participants have a few months to analyze the data and submit detection candidates, which are then compared with the sources originally injected in the data sets. (*Training* data sets with public source parameters are also provided to help participants tune and troubleshoot their codes.)

Three rounds of MLDCs had been completed at the time of this workshop, each spanning approximately 6 months. Challenge 1 [2, 3] was focused on establishing basic techniques to observe GWs from compact galactic binaries, intrinsically monochromatic, isolated or moderately interfering; as well as from the inspiral phase of bright, isolated, nonspinning massive-black-hole (MBH) binaries. Challenge 2 [4, 5] featured three considerably more complex data-analysis problems: a data set containing GW signals from approximately 26 million galactic binaries (again monochromatic) drawn from a randomized population-synthesis catalog; a data set (the ‘whole enchilada’) with a similar galactic-binary population, plus GW signals from an unknown number (between 4 and 6) of nonspinning-MBH binary and from five extreme-mass-ratio inspirals (EMRI); and five more data sets with single-EMRI signals.

The very steep increase in complexity introduced over a short time-scale with Challenge 2 and the need to consolidate analysis techniques (especially so for EMRIs) before moving to even more taxing challenges motivated the organization of Challenge 1B, a repeat of Challenge 1 with the addition of single-EMRI data sets. Challenge-1B data sets were distributed in the late summer 2007, with a deadline of December 2007 for entries. Ten collaborations submitted solutions. Highlights from this round include the range of techniques used, the participation of a number of new groups that successfully recovered signals from galactic binaries and MBH binaries, and the first convincing demonstration of EMRI detection *and* parameter estimation. Section 2 provides a brief summary of the entries; additional details about the work of individual collaborations are given elsewhere in this volume.

As we write (April 2008), Challenge 3 data sets have just been released, with entries due at the beginning of December 2008. Challenge 3 represents a definite step in the direction of more realistic source models (such as chirping galactic binaries and spinning-MBH binaries) and of new source classes (such as short-lived bursts and stochastic backgrounds). Section 3 describes the Challenge 3 data sets and waveform models in detail.

2. Report on Challenge 1B

Challenge 1B focused on three classes of GW sources, each tackled in a separate subchallenge: monochromatic galactic binaries, MBH binaries and EMRIs. The galactic-binary data sets (1B.1.1a–c and 1B.1.2–1.5) and the MBH-binary data sets (1B.2.1–2.2) had a duration of approximately 1 year (31 457 280 s, sampled at intervals of 15 s), while the EMRI data sets (1B.3.1–3.5) were twice as long (with the same sampling time). The challenge solutions submitted by the participants were assessed with the simple criteria adopted in previous rounds [3, 5]. Detector-response data were generated with the *best-fit* source parameters $\vec{\lambda}_{\text{sub}}$ submitted by the participants, using the same code previously employed to build the challenge data sets. These data were then compared to the detector response to the *true* waveforms, using as a figure of merit the *recovered* SNR

$$\text{SNR}(\vec{\lambda}_{\text{sub}}) = \frac{(A_{\text{true}}|A_{\text{sub}}) + (E_{\text{true}}|E_{\text{sub}})}{\sqrt{(A_{\text{sub}}|A_{\text{sub}}) + (E_{\text{sub}}|E_{\text{sub}})}}, \quad (1)$$

where A and E denote time series for the noise-orthogonal TDI observables $(2X - Y - Z)/3$ and $(Z - Y)/\sqrt{3}$ [6] and $(\cdot|\cdot)$ denotes the usual signal product weighted by instrument noise. We also quote the *correlation* $C = \text{SNR}/\text{SNR}_{\text{opt}}$, where $\text{SNR}_{\text{opt}} = \sqrt{(A_{\text{true}}|A_{\text{true}}) + (E_{\text{true}}|E_{\text{true}})}$ is the *optimal* SNR. For a perfect detection $C = 1$, but fluctuations $\sim 1/\text{SNR}_{\text{opt}}$ are expected because of instrument noise. When we examine parameter errors, these are defined simply as $\Delta\lambda^i = \lambda_{\text{sub}}^i - \lambda_{\text{true}}^i$; in some cases it makes sense to consider the *fractional* parameter errors $\Delta\lambda^i/\lambda^i = (\lambda_{\text{sub}}^i - \lambda_{\text{true}}^i)/\lambda_{\text{true}}^i$. In the remainder of this section we briefly discuss the entries submitted by participants; the technical notes accompanying the entries can be found at www.tapir.caltech.edu/~mldc/results1B/results.html.

2.1. Galactic binaries: Challenges 1B.1.X

Data sets 1B.1.1a–c and 1B.1.2–1.5 contained GW signals from monochromatic galactic binaries, in a variety of parameter ranges and source combinations. Seven parameters are required to fully characterize each such source: the amplitude \mathcal{A} , the (constant) frequency f , the ecliptic latitude and longitude β and λ , the inclination and polarization angles ι and ψ , and the initial phase ϕ_0 . Entries were submitted by five groups: *GSFC* (scientists at Goddard Space Flight Center), *IMPAN* (the Institute of Mathematics of the Polish Academy of Science

Table 1. Correlations for single-galactic-binary challenges 1B.1.1a–c. Asterisks denote entries corrected by maximizing the \mathcal{F} -statistic and using the resulting extrinsic parameters; these corrections are not reported where the frequency is well off, and the \mathcal{F} -statistic is merely fitting noise.

Group	1B.1.1a SNR _{opt} = 13.819	1B.1.1b SNR _{opt} = 24.629	1B.1.1c SNR _{opt} = 15.237
AEI	0.108 → 0.984*	0.922 → 0.996*	−0.190 → 0.989*
GSFC	0.992	0.807 → 0.814*	−0.138
IMPAN	0.988	0.981 → 0.997*	0.924 → 0.946*
MCMNJU	0.952 → 0.996*	0.906 → 0.994*	0.033
UIBBham	0.992	0.996	

and the Institute of Theoretical Physics at the University of Wrocław), *AEI* (the Albert Einstein Institute in Golm, Germany), *MCMNJU* (the Institute of Applied Mathematics of the Chinese Academy of Sciences and the Department of Astronomy of Nanjing University), *UIBBham* (the University of the Balearic Islands and the University of Birmingham). However, all groups except AEI concentrated only on a subset of the challenges.

Participants employed a fair range of techniques, in the same broad class as adopted for similar challenges in the past [3, 5]; however, new implementations and different technical solutions were pursued. *GSFC* used the X-Ray Spectral Fitting Package (XSPEC) [7] to fit templates to energy spectra. The package includes a Levenberg–Marquardt optimization algorithm, which was used to obtain an initial guess for the source parameters. A Markov-chain Monte Carlo (MCMC) routine, also available in XSPEC, was then used to converge to the best-fit source parameters. *IMPAN* set up a grid-based matched-filtering search with an optimized placement of templates on a hypercubic lattice. The \mathcal{F} -statistic [8] was used to reduce the search space from 7 to 3 parameters. A similar technique was adopted by *AEI* [9], in conjunction with a rigid-adiabatic model of detector response. *MCMNJU* used a genetic algorithm that optimized the \mathcal{F} -statistic; *UIBBham* implemented a MCMC search described in more detail in [10].

Each of data sets 1B.1.1a–c contained a GW signal from a single monochromatic binary (differing by frequency), with SNR \approx 13–25. Table 1 lists the correlations recovered by each collaboration. Some of the entries included close matches for the intrinsic parameters (f , θ and ϕ), but not for the remaining (extrinsic) parameters: this was due to the (relative) inaccuracy of the LISA response models used by the participants, or by mismatch between their definitions of the extrinsic parameters and the MLDC’s. In these cases we recomputed C by maximizing the \mathcal{F} -statistic [8] for the intrinsic parameters provided; the resulting C s are denoted by asterisks, and show that the intrinsic parameters were indeed recovered well enough to ensure solid detections. Table 2 lists parameter errors, and figure 1 shows where in the sky each collaboration placed the single binary of data set 1B.1.1a, compared to its true position, and to the Fisher-matrix $1\text{-}\sigma$ error contour.

Data set 1B.1.2 contained GW signals from 25 ‘verification’ binaries of known (i.e., disclosed) frequency and sky location. Five of them were taken from the list of observed binaries on Gijs Nelemans’ wiki [11], while the remaining 20 were placed randomly in the galaxy, varying their frequencies over a representative range. Table 3 lists the *global* correlations (computed for the combined signals of all reported and true binaries) recovered by the three groups that participated in this challenge. Just as it happened for Challenges 1B.1.1a–c, problems in assigning extrinsic parameters reduced the correlations. (Since the intrinsic parameters were provided to the participants, we did not perform \mathcal{F} -statistic-based




Figure 1. Sky positions reported by participants in Challenge 1B.1.1a, compared to the true location of the source. All the sky positions fall within the Fisher-matrix $1\text{-}\sigma$ contour, shown as a dotted line. (The spread of the reported sky positions is due to both the difference in noise realizations between the Synthetic LISA and LISA Simulator versions of data set 1B.1.1a, and the systematic errors of the searches.)

Table 2. Parameter errors for Challenges 1B.1.1a–c. All angles are expressed in radians.

Group	$\Delta\beta$	$\Delta\lambda$	Δf (nHz)	$\Delta\psi$	Δt	$\Delta\varphi$	$\Delta\mathcal{A}$ (10^{-23})
Challenge 1B.1.1a ($f_{\text{true}} = 1.060$ mHz)							
AEI	-0.032	-0.120	-2.43	0.217	-0.454	1.17	1.22
GSFC	-0.004	-0.071	-1.81	0.708	0.252	1.33	1.20
IMPAN	-0.031	0.018	2.13	0.454	0.212	-1.06	1.25
MCMNJU	-0.017	-0.042	-0.53	0.662	0.426	-1.57	2.34
UIBBham	0.005	-0.079	-1.51	0.708	0.173	-1.32	0.65
Challenge 1B.1.1b ($f_{\text{true}} = 2.904$ mHz)							
AEI	-0.056	-0.0090	0.95	-1.050	0.283	1.63	-0.066
GSFC	-0.462	0.0606	-30.9	2.560	0.182	0.52	-0.024
IMPAN	0.020	0.0007	0.85	0.333	0.339	-0.60	0.713
MCMNJU	-0.067	-0.0063	2.07	-0.732	-0.064	0.84	-0.223
UIBBham	-0.044	-0.0082	1.78	-0.636	0.043	1.13	-0.029
Challenge 1B.1.1c ($f_{\text{true}} = 9.943$ mHz)							
AEI	-0.026	0.0053	1.84	-0.499	-1.120	3.02	0.124
GSFC	-0.452	-1.48	140	1.820	-0.471	-0.66	-0.695
IMPAN	-0.016	0.0248	3.72	-1.510	-0.197	2.68	0.478
MCMNJU	-0.555	-0.3680	359	-1.590	-0.250	-0.94	-0.532

adjustments, which would amount to solving the entire problem. As a result, the low C values of table 3 may be symptomatic only of extrinsic-parameter systematics.)

Data set 1B.1.3 contained GW signals from 20 unknown binaries distributed across the LISA band, well separated in frequencies. Unfortunately, a bug in the random generation of source parameters caused all SNRs to be too small for detection (all were below 1). Happily, no

Table 3. Correlations, number of recovered sources, and number of false positives for Challenges 1B.1.2, 1B.1.4 and 1B.1.5. Correlations computed after correcting extrinsic parameters using the \mathcal{F} -statistic are denoted with asterisks.

Group	C	# recovered
1B.1.2 (SNR _{opt} = 634.918, 25 sources)		
AEI	-0.822	25
GSFC	0.006	25
MCMNJU	0.267	25
1B.1.4 (SNR _{opt} = 340.233, 51 sources)		
AEI	0.774 → 0.966*	13 (2 false positives)
GSFC	0.003 → 0.282*	6 (1 false positives)
1B.1.5 (SNR _{opt} = 273.206, 44 sources)		
AEI	0.453 → 0.929*	3

participating group reported a positive detection, consistent with the correct behavior expected of search algorithms.

Challenge 1B.1.4 was meant to test search algorithms in the presence of mild source confusion. Fifty-one sources were spread across a band of 15 μHz beginning at 3 mHz, with an average density of 0.108 sources per frequency bin. By contrast, Challenge 1B.1.5 tested algorithms in the presence of a higher level of source confusion, comparable to that expected from our galaxy. Forty-four sources were spread across a band of 3 μHz centered at 3 mHz, with an average density of 0.465 sources per frequency bin. Table 3 lists the global correlations and the number of sources recovered by the participating groups, as well as the number of *false positives*, defined here as reported sources farther than one frequency bin (1/year) from any true source, or with \mathcal{F} -statistic-adjusted correlation less than 0.7 with all true sources within a frequency bin. The top panel of figure 2 shows the combined A and E spectral amplitude for the GW signals in data set 1B.1.4, together with the residual after subtracting the signal model submitted by *AEI*. The bottom panel shows the same subtraction after extrinsic parameters have been recomputed for this entry by maximizing the \mathcal{F} -statistic.

Altogether, it must be said that Challenge 2 provided a more forceful demonstration of LISA's science objectives for galactic binaries [5]; but Challenge 1B was still very useful for new groups to start implementing search methods, and for established groups to continue tuning them. The extrinsic-parameter reporting errors seen here are easy to commit, because these parameters are very sensitive to the modeling of the LISA response, and because their definitions are somewhat conventional; but these errors have little bearing on detection confidence. To avoid such problems in the future, we plan to provide a web tool to check the recovered SNR against the challenge data sets using the fiducial MLDC waveform-generation code.

2.2. Massive black hole binary systems: Challenges 1B.2.X

Each of data sets 1B.2.1 and 1B.2.2 contained a loud GW signal from a single MBH binary embedded in instrument noise. Gravitational radiation was modeled as the *restricted* waveform for spinless point masses moving on an adiabatic sequence of circular orbits, evolving according to 2PN energy-balance equations [4, section 4.4]. (See also section 3.2 for the new waveform features being introduced for Challenge 3.) Nine parameters are needed to describe each source: the two masses m_1 and m_2 , the time of coalescence t_c , the sky-position angles β and λ , the luminosity distance D_L , the orbital-inclination and GW-polarization angles ι and ψ , and the initial orbital phase φ_0 . The binaries in both data sets had masses drawn from the




Figure 2. Combined A and E spectral amplitude for the GW signals in data set 1B.1.4, before and after subtracting the AEI signal model. In the bottom panel, the extrinsic parameters for all binaries in the model were adjusted by maximizing the \mathcal{F} -statistic.

(This figure is in colour only in the electronic version)

same ranges ($m_1 = 1\text{--}5 \times 10^6 M_\odot$, $m_1/m_2 = 1\text{--}4$), but were distinguished by the times of coalescence $t_c = 6 \pm 1$ months for Challenge 1B.2.1 and 400 ± 40 days (past the end of the data set) for Challenge 1B.2.2; the SNR_{opt} for the sources were chosen to be $\simeq 500$ and $\simeq 80$, respectively.

Two groups submitted entries: *JPL* (a collaboration between researchers at Caltech and at the Jet Propulsion Laboratory) employed a three-step hierarchical strategy combining a time–frequency track-search analysis, a template-bank matched-filtering search, and a final MCMC stage to evaluate the posterior probability densities of source parameters for data sets 1B.2.1 and 1B.2.2. *Cardiff* (a collaboration based at that university) used a stochastic-template-bank matched-filtering search [12] to analyze data set 1B.2.1.

Table 4 summarizes these entries. While the recovered SNRs are very close to SNR_{opt} , which indicates detections of very high confidence, there are large discrepancies in the sky-position angles for Challenge 1B.2.1, where both the *JPL* and *Cardiff* searches converged on secondary likelihood maxima, very close in height to the true mode (as shown by the recovered SNR), but quite distant in parameter space. In fact, the *JPL* result places the source almost at the antipodal sky position, even if $\text{SNR} \simeq \text{SNR}_{\text{opt}}$ to better than one part in a thousand.

This is a true global degeneracy, which does not appear in local Fisher-matrix analyses (another example of why mock-data endeavors are useful!), and which may indicate the need, in EM searches of counterparts to LISA binary-MBH detections, to examine unconnected regions of the sky. It may however be premature to make such an inference, since the degeneracy could be broken by spin effects (now included in Challenge 3 waveforms) and higher waveform harmonics. In addition, EM-counterpart searches would require a data-analysis system capable of determining the sky position of MBH binaries a few days in advance of their merger (corresponding to the interval between data dumps from LISA to the ground), whereas data set 1B.2.1 included the inspiral waveform all the way to the approximate merger frequency. Therefore this challenge was *not* aimed directly at establishing the feasibility of sky-position determination for EM-counterpart searches. In this context, it is however worth

Table 4. Relative/absolute errors for the MBH binaries in Challenges 1B.2.1 and 1B.2.2. All angles are given in radians. Error estimates have been adjusted to account for two *perfect* symmetries of the waveforms with respect to source parameters: $\psi \rightarrow \psi + \pi$, and the simultaneous transformation $\psi \rightarrow \psi + \pi/2$, $\varphi_0 \rightarrow \varphi_0 - \pi/2$.

Group	1B.2.1 (SNR _{opt} = 531.84)		1B.2.2 (SNR _{opt} = 80.67)
	JPL	Cardiff	JPL
SNR	531.57	511.78	79.86
$\Delta m_1/m_1$	5.991×10^{-3}	0.108	0.122
$\Delta m_2/m_2$	-5.252×10^{-3}	-0.111	-0.134
Δt_c [s]	206.1	-541.8	-2688.2
$\Delta D_L/D_L$	-0.139	-1.438	4.781×10^{-3}
$\Delta \beta$	2.429	1.374	5.862×10^{-3}
$\Delta \lambda$	3.133	0.548	-1.461×10^{-2}
$\Delta \iota$	0.713	0.678	-6.955×10^{-2}
$\Delta \psi$	-0.564	1.448	-4.878×10^{-2}
$\Delta \varphi_0$	-2.846	-2.389	1.293×10^{-2}

pointing out that the entries provided very accurate determinations of the times of coalescence, corresponding to time windows of a few minutes (for Challenge 1B.2.1) and about 45 min (for 1B.2.2).

If we compare the errors of table 4 with the parameter-determination accuracies predicted in the Fisher-matrix formalism, we see that the *JPL* estimates for m_1 and m_2 fall within the $2\text{-}\sigma$ contour for data set 1B.2.1 (notwithstanding the problem with sky position), and near the $1\text{-}\sigma$ contour for 1B.2.2. For data set 1B.2.2, *JPL*'s Δt_c , ΔD_L , $\Delta \beta$, and $\Delta \varphi_0$, are also close to $1\text{-}\sigma$, and $\Delta \lambda$ is $\sim 2.2\sigma$; the errors in ι and ψ are all tens of σ s (but the Fisher-matrix formalism is not always reliable for extrinsic parameters). Altogether, we conclude that the *JPL* search essentially achieves the theoretical limits of parameter extraction for data set 1B.2.2, while it does so for an important subset of parameters for data set 1B.2.1.

2.3. Extreme mass ratio inspirals: Challenges 1B.3.X

Although Challenge 2 saw a few successful detections of EMRI signals [5], data sets 1B.3.1–3.5 represented the first real testbed for the search algorithms developed for this critical source class. Each data set contained a GW signal from a single EMRI embedded in instrument noise, with SNR_{opt} between $\simeq 55$ and $\simeq 135$, and source parameters chosen randomly as described in [4]. Fourteen parameters are needed to describe each EMRI source [4]: the ecliptic latitude and longitude β and λ and the luminosity distance D_L ; the central-BH and compact-object masses M and μ ; the magnitude a and orientation angles θ_K , ϕ_K of the central-BH spin; the initial radial orbital frequency ν_0 and eccentricity e_0 ; and three angles $\tilde{\gamma}_0$, α_0 and λ describing the initial orientation of the orbit.

Entries were received from three groups: *BBGP* (a collaboration of scientists at the AEI, Cambridge, and the University of Southampton); *EtfAG* (AEI, Northwestern, and Cambridge); and *MT* (Montana State University). *EtfAG* employed a time–frequency technique [13], whereas *BBGP* [14] and *MT* [15] developed coherent approaches based on Monte Carlo techniques, differing in their implementation. The entries were assessed as discussed at the beginning of this section, although the *EtfAG* time–frequency analysis cannot determine the extrinsic parameters, so their recovered SNR and correlation could not be computed. Tables 5 and 6 summarize all results.

Both the time–frequency and coherent approaches succeeded in detecting these relatively strong EMRI signals and in constraining their parameters (with especially remarkable accuracy

Table 5. Overlaps and recovered SNRs for TDI observables A , E and combined recovered SNR for data sets 1B.3.1–3.5.

Group	C_A	SNR_A	C_E	SNR_E	Total SNR
1B.3.1 ($\text{SNR}_{\text{opt}} = 123.7$)					
BBGP	0.57	51.0	0.58	51.6	72.5
MT	0.998	86.1	0.997	88.3	123.4
1B.3.2 ($\text{SNR}_{\text{opt}} = 133.5$)					
BBGP	0.07	6.6	0.18	18.2	17.6
BBGP ^a	0.39	37.6	0.41	39.8	54.7
MT	0.54	49.5	0.54	50.8	70.9
1B.3.3 ($\text{SNR}_{\text{opt}} = 81.0$)					
BBGP	−0.06	−4.2	−0.0003	−0.05	−3.0
BBGP ^{a,b}	−0.2	−11.5	−0.32	−19.0	−21.5
MT	0.38	22.0	0.35	20.9	30.4
1B.3.4 ($\text{SNR}_{\text{opt}} = 104.5$)					
BBGP ^b	0.0007	2.1	−0.0002	−0.8	2.1
BBGP ^c	0.16	13.9	0.04	6.7	14.6
1B.3.5 ($\text{SNR}_{\text{opt}} = 57.6$)					
BBGP	0.09	3.4	0.1	4.2	5.3

^a C and SNR after correcting the sign of β , lost on input to the MLDC webform.

^b The *BBGP* SNRs can be negative because *BBGP* maximized likelihood analytically over amplitude, which makes SNR sign-insensitive (a minus sign corresponds to a change of π in the phase of the dominant harmonic). This degeneracy is broken when all the harmonics are found correctly.

^c C and SNR after correcting phases at $t = 0$, to account for a *BBGP* bug.

for data set 1B.3.1), although not all groups analyzed all data sets, and the performance of the same search pipeline varied across them. Challenge participants report that in some cases this was due to a lack of time for extended computations before the challenge deadline, so some parameter set were submitted as ‘best fits’ although they were clearly understood to be secondary likelihood maxima. Thus, this early development work indicates that the main challenge for (isolated) EMRI analyses is the very complex structure of the likelihood surface in source-parameter space, which features a number of secondary maxima of similar height, even more so than for MBH binaries.

We caution the reader that it would be inappropriate at this time to draw general conclusions about the relative merits of search methods and about the expected science payoff of LISA EMRI astronomy: it is not known how these techniques scale as the SNR decreases and in situations where EMRI signals overlap with each other and are affected by galactic confusion noise. The first two complications will be addressed in Challenge 3.

3. Synopsis of Challenge 3

The third round of the MLDCs consists of five challenges (3.1–3.5). Data sets 3.1–3.3 consist of approximately 2 years of data (2^{22} samples at a cadence of 15 s) for time-delay interferometry (TDI) observables X , Y and Z . These data sets are released both as time series of equivalent strain generated by the LISA Simulator [18] and as time series of fractional frequency fluctuations generated by Synthetic LISA [19]; see [4, p S556] for the conversion

Table 6. Errors for a subset of EMRI parameters in Challenges 1B.3.1–3.5. ‘ $\Delta x/[x]$ ’ denote fractional errors relative to the physical or prior range of the parameter. (This shows the amount of information about the parameters that is gained with the search.) Large errors correspond to searches converging to secondary maxima of the likelihood; the true global maximum was found only by *MT* in Challenge 1B.3.1.

Group	$\frac{\Delta\beta}{[\beta]}$	$\frac{\Delta\lambda}{[\lambda]}$	$\frac{\Delta\theta_K}{[\theta_K]}$	$\frac{\Delta\phi_K}{[\phi_K]}$	$\frac{\Delta a}{[a]}$	$\frac{\Delta\mu}{[\mu]}$	$\frac{\Delta M}{[M]}$	$\frac{\Delta v_0}{v_0}$	$\frac{\Delta e_0}{0.15}$	$\frac{\Delta\lambda_{SL}}{[\lambda_{SL}]}$
Challenge 1B.3.1										
BBGP	-0.03	-0.0059	-0.14	0.053	0.31	-0.20	-0.84	0.026	0.37	-0.022
EtfAG	0.019	-0.0045	0.56	0.33	0.16	-0.11	-0.27	-9.3×10^{-5}	0.17	0.078
MT	0.0058	0.0027	4.4×10^{-4}	0.0051	-0.0022	0.0065	0.014	3.2×10^{-6}	-0.0085	-0.0020
Challenge 1B.3.2										
BBGP	-0.16	-0.43	0.46	-0.33	-0.0088	-0.0040	0.016	1.4×10^{-4}	-0.010	-0.0013
EtfAG	-0.014	0.0042	0.97	-0.36	0.0043	-0.046	-0.069	-6.5×10^{-5}	0.041	0.0041
MT	0.0040	-0.0086	0.79	0.41	0.093	-0.064	0.35	-0.035	0.068	0.092
Challenge 1B.3.3										
BBGP	0.091	0.50	-0.23	0.045	-0.32	-0.49	-0.029	6.1×10^{-4}	0.019	0.054
EtfAG	-0.01	-0.004	0.49	-0.34	0.0073	-0.059	-0.061	-7.8×10^{-5}	0.038	0.0061
MT	0.045	-0.019	-0.1	0.077	-0.066	0.13	0.59	3.6×10^{-4}	-0.33	0.010
Challenge 1B.3.4										
BBGP	-0.57	-0.37	0.37	-0.31	-0.025	0.020	-0.88	0.066	0.065	-0.16
EtfAG	-0.56	0.49	0.56	-0.34	0.059	0.12	0.04	2.8×10^{-4}	-0.039	0.0040
Challenge 1B.3.5										
BBGP	-0.48	-0.14	-0.35	0.1	-0.094	-0.094	0.55	-0.0021	-0.017	-0.060
EtfAG	-0.58	0.46	0.27	-0.084	0.20	-0.7	0.83	-0.066	0.066	0.27

between the two. Indeed (with a few exceptions, described below, for 3.4 and 3.5), the Challenge-3 data sets are built using the ‘pseudo-LISA’ model of Challenges 1 and 2: the orbits of the LISA spacecraft are e^2 -accurate Keplerian ellipses with conventional orientations and time offsets; *modified* TDI (a.k.a. TDI 1.5) expressions are used for the observables; and Gaussian, stationary instrument noise is included from six proof masses and six optical benches with known noise levels that are identical across each set of six²². See [4] for details.

- *Data set 3.1* contains a galactic GW foreground from ~ 60 million compact binary systems. This data set is a direct descendant of Challenge 2.1, but it improves on the realism of the latter by including both detached and interacting binaries with intrinsic frequency drifts (either positive or negative). Section 3.1 gives details about the binary waveform models,

²² The six proof-mass noises are uncorrelated and white in acceleration, with one-sided power spectral density (PSD)

$$S_{\text{acc}}^{1/2}(f) = 3 \times 10^{-15} [1 + (10^{-4} \text{ Hz}/f)^2]^{1/2} \text{ m s}^{-2} \text{ Hz}^{-1/2};$$

the six optical-path noises are uncorrelated and white in phase with PSD

$$S_{\text{opt}}^{1/2}(f) = 20 \times 10^{-12} \text{ m Hz}^{-1/2};$$

the conversion to Synthetic LISA’s dimensionless fractional frequency fluctuations is described on [19, p 6]; the values actually used in the MLDCs are

$$S_{\text{acc}}(f) = 2.5 \times 10^{-48} (f/\text{Hz})^{-2} [1 + (10^{-4} \text{ Hz}/f)^2] \text{ Hz}^{-1};$$

$$S_{\text{opt}}(f) = 1.8 \times 10^{-37} (f/\text{Hz})^2 \text{ Hz}^{-1}.$$

about their implementation in the LISAtools suite [16] and about the generation of the galactic population.

- *Data set 3.2* contains GW signals from 4–6 binaries of spinning MBHs, on top of a confusion galactic-binary background. This data set improves on the realism of Challenges 1.2.1–2 and 2.2 by modeling the orbital precession (and ensuing GW modulations) due to spin–orbit and spin–spin interactions. Section 3.2 gives details about the MBH-binary waveforms.

Because this challenge focuses on the effects of spins rather than on the joint search for MBH signals and for the brightest galactic binaries, the background is already *partially subtracted*—it is generated from the population of detached binaries used for Challenge 3.1, withholding all signals with $\text{SNR} > 5$.

- *Data set 3.3* contains five GW signals from EMRIs. As in Challenges 1.3.1–5, EMRI waveforms are modeled with Barack and Cutler’s ‘analytic kludge’ waveforms [17]; this challenge introduces the complication of detecting five such signals with lower SNRs, and *in the same data set*. By contrast, galactic confusion is not included. See section 3.3 for details.

Challenges 3.4 and 3.5 address the detection of two GW sources that are *new* to the MLDCs, and that have (respectively) bursty and stochastic characters: thus, these searches require an accurate characterization of instrument noise, which in reality will not be available *a priori*, but will be obtained from the LISA measurements themselves. To model this problem, in data sets 3.4 and 3.5 the levels of the six + six secondary noises have been independently randomized by $\pm 20\%$; the noises are however still uncorrelated. In addition, these data sets contain time series for all 12 ‘raw’ LISA phase measurements y_{ijk} and z_{ijk} [19], so that challenge participants may now build additional TDI observables to help characterize instrument noise. The phase measurements *do* include laser phase noise, because otherwise they would convey extra information unavailable from the real LISA; but laser noise is reduced in level to ~ 10 times the secondary noise at 1 mHz, so that it can be canceled relatively easily with TDI 1.5 implemented with moderate timing precision. To wit:

- *Data set 3.4* consists of 2^{21} samples at a cadence of 1 s (~ 24 days altogether), and it contains GW burst signals from cosmic string cusps, occurring as a Poissonian random process throughout the data set, with an expectation value of five events. Details about the waveforms are given in section 3.4. The data set is provided only as fractional frequency fluctuations generated by Synthetic LISA.
- *Data set 3.5* consists of 2^{20} samples at a cadence of 2 s (again ~ 24 days), and it contains a stochastic GW background, which is isotropic, unpolarized, Gaussian and stationary; its spectrum grows at low frequencies as $1/f^3$, and its magnitude is set to a few times the secondary noise over a broad range of frequencies. Details about the synthesis of the background and the simpler model of the LISA orbits used for this challenge are given in section 3.5. The data set is provided as fractional frequency fluctuations generated by Synthetic LISA and by the new simulator LISACode [20], recently integrated into the LISAtools suite [16]; thus, cross checks are possible between the two simulators.

LISACode [20] was developed at APC-Paris with the purpose of accurately mapping the impact of the different LISA subsystems on its science observations, and of bridging the gap between the basic principles of the LISA measurement and a future, more sophisticated end-to-end simulator. Thus, LISACode includes realistic representations of most of the ingredients that will influence LISA’s sensitivity (such as orbits, instrument noise, ultra-stable-oscillator time stamps, phasemeter response functions), internal waveform generators for several kinds of sources (monochromatic and chirping binaries, stochastic backgrounds, etc), as well as

the possibility of building various TDI combinations. Many user-defined parameters make it possible to study the impact of different LISA configurations on its sensitivity. LISACode's conventions follow closely those of the MLDCs and of Synthetic LISA.

All the Challenge-3 data sets can be downloaded at astrograves.nasa.gov/docs/mldc/round3/datasets.html, encoded in lisaXML [2], an XML-based format that can be displayed directly in modern web browsers, and handled easily in C/C++, Python and MATLAB with the LISAtools I/O libraries [16]. Each data set is released in the blind challenge version and in a training version that includes the source parameters used to generate it. Additional training data sets can be generated easily with the LISAtools suite²³.

The remainder of this section describes the GW signal models adopted for each data set. See [4] for the conversion of the GW polarizations in source frame (given here) to the LISA frame. Table 7 is a glossary of source parameters with their symbols and lisaXML descriptors, while table 8 is a summary of the GW content of each data set along with the ranges used to choose source parameters randomly.

3.1. Chirping galactic binaries

Data set 3.1 contains GWs from a population of $\sim 26 \times 10^6$ detached and $\sim 34 \times 10^6$ interacting galactic binaries. Each binary is modeled as a system of two point masses m_1 and m_2 in circular orbit with linearly increasing or decreasing frequency (depending on whether gravitational radiation or equilibrium mass transfer is dominant). The polarization amplitudes at the solar-system barycenter, expressed in the source frame, are given by

$$\begin{aligned} h_+^S(t) &= \mathcal{A}(1 + \cos^2 \iota) \cos[2\pi(ft + \dot{f}t^2/2) + \phi_0], \\ h_\times^S(t) &= -2\mathcal{A}(\cos \iota) \sin[2\pi(ft + \dot{f}t^2/2) + \phi_0], \end{aligned} \quad (2)$$

where the amplitude is derived from the physical parameters of the source as $\mathcal{A} = (2\mu/D_L)(\pi Mf)^{2/3}$, with $M = m_1 + m_2$ the total mass, $\mu = m_1 m_2 / M$ the reduced mass, and D_L the distance; \dot{f} is the (constant) frequency derivative, and ϕ_0 is the phase at $t = 0$.

Since it would be unfeasible to process millions of barycentric binary waveforms individually through the LISA simulators to compute the TDI-observable time series, we adopt a fast frequency-domain method [21] that rewrites the LISA phase measurements as the fast-slow decomposition

$$y_{ij}(t) = C(t) \cos(2\pi f_0 t) + S(t) \sin(2\pi f_0 t); \quad (3)$$

the functions $C(t)$ and $S(t)$ describe slowly varying effects such as the rotation of the LISA arms, the Doppler shift induced by orbital motion, and the intrinsic frequency evolution of the source. These 'slow' terms can be sampled very sparsely and Fourier-transformed numerically, while the 'fast' sine and cosine terms can be Fourier-transformed analytically. The results are then convolved to produce the LISA phase measurements, and these are assembled into the desired TDI variables. This algorithm is three-to-four orders of magnitude faster than the time-domain LISA simulators, although it effectively approximates LISA as a rigidly rotating triangle with equal and constant armlengths. See [21] for full details, and directory MLDCwaveforms/galaxy3 in LISAtools for the source code.

The starting point for each realization of data set 3.1 are two large catalogs provided by Gijs Nelemans (files MLDCwaveforms/galaxy3/Data/AMCVn_GWR_MLDC.dat and

²³ After installing LISAtools following the instructions at code.google.com/p/lisatools/wiki/Install, generating a training set is as simple as running (say, for Challenge 3.1)

```
MLDCpipelines2/bin/challenge3.py -T -R 3.1.
```

Table 7. Source parameters in Challenge 3.

Parameter	Symbol	Standard parameter name (lisaXML descriptor)	Standard unit (lisaXML descr.)
<i>Common parameters</i>			
Ecliptic latitude	β	EclipticLatitude	Radian
Ecliptic longitude	λ	EclipticLongitude	Radian
Polarization angle	ψ	Polarization	Radian
Inclination	ι	Inclination	Radian
Luminosity distance ^a	D_L	Distance	Parsec
<i>Galactic binaries</i>			
Amplitude ^b	A	Amplitude	1 (GW strain)
Frequency	f	Frequency	Hertz
Frequency derivative	\dot{f}	FrequencyDerivative	Hertz/second
Initial GW phase	ϕ_0	InitialPhase	Radian
<i>Spinning massive black-hole binaries</i>			
Masses of component MBHs	m_1, m_2	Mass1, Mass2	SolarMass
Magnitude of spins S_1, S_2	a_1, a_2	Spin1, Spin2	MassSquared
Initial orientation of spin S_1	θ_{S1}, ϕ_{S1}	PolarAngleOfSpin1 AzimuthalAngleOfSpin1	Radian Radian
Initial orientation of spin S_2	θ_{S2}, ϕ_{S2}	... <i>likewise</i>	
Time to coalescence	T_c	CoalescenceTime	Second
Phase at coalescence	Φ_c	PhaseAtCoalescence	Radian
Initial orientation of orbital momentum	θ_L, ϕ_L	InitialPolarAngleL InitialAzimuthalAngleL	Radian Radian
<i>EMRIs: see table 5 of [4]</i>			
<i>Cosmic string cusp bursts</i>			
Amplitude ^b (Fourier)	A	Amplitude	Hertz ^{1/3}
Central time of arrival	t_C	CentralTime	Second
Maximum frequency ^c	f_{\max}	MaximumFrequency	Hertz
<i>Isotropic stochastic background</i>			
PSD ^{b,d} at 1 Hz	S_h	PowerSpectralDensity	(f/Hz) ⁻³ /Hz

^a We do not deal explicitly with the redshifting of sources at cosmological distances, so D_L is a *luminosity* distance, and all masses and frequencies are measured at the solar-system barycenter and red/blue-shifted by factors $(1+z)^{\pm 1}$ with respect to those measured locally near the sources.

^b Replaces D_L for galactic binaries, cosmic-string-cusp bursts and stochastic-background pseudosources.

^c Effectively replaces ι for cosmic-string-cusp bursts.

^d Note also that $S_h = S_h^{\text{rot}}/384$; ψ is set to 0, and ι not used.

dwd_GWR_MLDC.dat in the LISAtools installation), which contain the parameters of 26.1×10^6 detached and 34.2×10^6 interacting systems produced by the population synthesis codes described in [22, 23]. Figure 3 shows the distribution of the binaries in the catalogs over f and \dot{f} . Recent work by Roelofs, Nelemans and Groot [24] suggests that the model in [23] overpredicts the number of (AM CVn) interacting systems by a factor of 5–10, but we did not implement this correction for Challenge 3.

The parameters of each binary in the catalogs are modified by randomly tweaking f by $\pm 1\%$, A by $\pm 10\%$, β and λ by $\pm 0.5^\circ$, and by randomly assigning ψ , ι and ϕ_0 (\dot{f} is computed from the catalog's binary-period derivative and from the tweaked f). These random perturbations are large enough to render the original population files useless as answer keys,

Table 8. Summary of data-set content and source-parameter selection in Challenge 3. Parameters are sampled randomly from uniform distributions across the ranges given below, and all angles (including spin and orbital-angular-momentum directions for MBH binaries) across the entire appropriate ranges. Source distances are set from individual-source SNRs, which are drawn randomly from the ranges specified below (where ‘SNR’ refers to the multiple-TDI-observable SNR approximated as $\sqrt{2} \times \max\{\text{SNR}_X, \text{SNR}_Y, \text{SNR}_Z\}$). The MBH time of coalescence t_c and the cosmic-string-cusp burst central time t_C are given relative to the beginning of the relevant data sets.

Data set	Sources	Parameters
3.1	<i>Galactic-binary background</i> plus 20 <i>verification binaries</i>	Randomized population (see section 3.1) $\sim 34 \times 10^6$ interacting, $\sim 26 \times 10^6$ detached Known parameters (see section 3.1)
3.2	4–6 <i>MBH binaries</i> ... including ... and 2–4 chosen from plus <i>galactic confusion</i>	for each: $m_1 = 1\text{--}5 \times 10^6 M_\odot$, $m_1/m_2 = 1\text{--}4$, $a_1/m_1 = 0\text{--}1$, $a_2/m_2 = 0\text{--}1$ MBH ₁ : $t_c = 90 \pm 30$ days, SNR ~ 2000 MBH ₂ : $t_c = 765 \pm 15$ days, SNR ~ 20 MBH ₃ : $t_c = 450 \pm 270$ days, SNR ~ 1000 MBH ₄ : $t_c = 450 \pm 270$ days, SNR ~ 200 MBH ₅ : $t_c = 540 \pm 45$ days, SNR ~ 100 MBH ₆ : $t_c = 825 \pm 15$ days, SNR ~ 10 randomized population with approx. SNR < 5 $\sim 26 \times 10^6$ binaries; no verification
3.3	5 <i>EMRIs</i> ... including	for each: $\mu = 9.5\text{--}10.5 M_\odot$, $S = 0.5\text{--}0.7 M^2$, time at plunge = $2^{21}\text{--}2^{22} \times 15$ s, ecc. at plunge = 0.15–0.25, SNR = 10–50 emri ₁ : $M = 0.95\text{--}1.05 \times 10^7 M_\odot$ emri ₂ and emri ₃ : $M = 4.75\text{--}5.25 \times 10^6 M_\odot$ emri ₄ and emri ₅ : $M = 0.95\text{--}1.05 \times 10^6 M_\odot$
3.4	n <i>Cosmic-string-cusp bursts</i>	(with n Poisson-distributed with mean 5) $f_{\text{max}} = 10^{-3}\text{--}1$ Hz, $t_C = 0\text{--}2^{21}$ s, SNR = 10–100 all instrument noise levels randomized $\pm 20\%$
3.5	<i>Isotropic stochastic background</i>	2×192 incoherent h_+ and h_\times sources over sky $S_h^{\text{tot}} = 0.7\text{--}1.3 \times 10^{-47} (f/\text{Hz})^{-3} \text{ Hz}^{-1}$ all instrument noise levels randomized $\pm 20\%$

but small enough to preserve the overall parameter distributions. Binaries with approximate single-Michelson SNR > 10 are regarded as ‘bright’ and listed in a separate table in the challenge keys. Data set 3.1 also includes 20 verification binaries of known parameters (specified in LISAtools file MLDCwaveforms/galaxy3/Data/Verification.dat as rows of f , \dot{f} , β , λ , A).

3.2. Spinning MBH binaries

The spinning-MBH-binary GW signals of data set 3.2 are modeled as restricted waveforms (no higher harmonics) from 2PN circular adiabatic inspirals, with uncoupled orbital frequency evolution and spin and orbital precession. Both the orbital phase and frequency are computed as explicit functions of time, corresponding to $T3$ waveforms in the classification of [25]:




Figure 3. Histogram of the density of galactic binaries in the Nelemans catalogs, binned by $\log_{10} f$ and $\log_{10} |\dot{f}|$.

$$M\omega = \frac{1}{8}\tau^{-3/8} \left[1 + \left(\frac{743}{2688} + \frac{11}{32}\eta \right) \tau^{-1/4} - \frac{3}{10} \left(\pi - \frac{\beta}{4} \right) \tau^{-3/8} + \left(\frac{1855099}{14450688} + \frac{56975}{258048}\eta + \frac{371}{2048}\eta^2 - \frac{3}{64}\sigma \right) \tau^{-1/2} \right], \quad (4)$$

where $M = m_1 + m_2$ is the total mass, $\eta = m_1 m_2 / M^2 \equiv \mu / M$ is the symmetric mass ratio, and

$$\tau = \frac{\eta}{5M} (T_c - t), \quad (5)$$

$$\beta = \frac{1}{12} \sum_{i=1,2} \left[\chi_i (\hat{\mathbf{L}}_N \cdot \hat{\mathbf{S}}_i) \left(113 \frac{m_i^2}{M^2} + 75\eta \right) \right], \quad (6)$$

$$\sigma = -\frac{1}{48} \eta \chi_1 \chi_2 [247(\hat{\mathbf{S}}_1 \cdot \hat{\mathbf{S}}_2) - 721(\hat{\mathbf{L}}_N \cdot \hat{\mathbf{S}}_1)(\hat{\mathbf{L}}_N \cdot \hat{\mathbf{S}}_2)]. \quad (7)$$

Here $\hat{\mathbf{L}}_N$, $\hat{\mathbf{S}}_1$ and $\hat{\mathbf{S}}_2$ are the unit vectors along the leading-order angular orbital momentum and the MBH spins. The *intrinsic* orbital phase is

$$\Phi_{\text{orb}} = \Phi_C - \frac{\tau^{5/8}}{\eta} \left[1 + \left(\frac{3715}{8064} + \frac{55}{96}\eta \right) \tau^{-1/4} - \frac{3}{16} (4\pi - \beta) + \left(\frac{9275495}{14450688} + \frac{284875}{258048}\eta + \frac{1855}{2048}\eta^2 - \frac{15}{64}\sigma \right) \tau^{-1/2} \right]; \quad (8)$$

however, because the spin-orbit coupling causes the orbital angular momentum to precess around the total angular momentum, the phase that enters the gravitational waveforms contains the additional correction [26]:

$$\dot{\Phi} = \omega + \frac{(\hat{\mathbf{L}}_N \cdot \hat{\mathbf{n}})[\hat{\mathbf{L}}_N \times \hat{\mathbf{n}}] \cdot \hat{\mathbf{L}}_N}{1 - (\hat{\mathbf{L}}_N \cdot \hat{\mathbf{n}})^2} \equiv \omega + \delta\dot{\Phi}, \quad (9)$$

where $\hat{\mathbf{n}}$ is direction to the source. The constant of integration in this equation can be redefined so that $\delta\Phi = 0$ at $t = 0$. The equations of precession for $\hat{\mathbf{L}}_N$, $\hat{\mathbf{S}}_1$ and $\hat{\mathbf{S}}_2$ are given by (2.9)–(2.11) in [27]. In the source frame the gravitational polarizations are then given (with respect to a time-varying polarization basis) by

$$\begin{aligned} h_+ &= -\frac{2\mu}{D}(1 + \cos^2 i)(M\omega)^{2/3} \cos 2\Phi, \\ h_\times &= \frac{4\mu}{D} \cos i (M\omega)^{2/3} \sin 2\Phi, \end{aligned} \quad (10)$$

where $\cos i = (\hat{\mathbf{L}}_N \cdot \hat{\mathbf{n}})$. The polarizations h_+^S and h_\times^S in a fixed source frame are obtained by way of a rotation by the instantaneous polarization angle

$$\tan \psi = \frac{\sin \beta \cos(\lambda - \phi_L) \sin \theta_L - \cos \theta_L \cos \beta}{\cos \beta \sin(\lambda - \phi_L)}, \quad (11)$$

(where θ_L and ϕ_L define the direction of \mathbf{L}_N) yielding

$$h_+^S = -h_+ \cos 2\psi - h_\times \sin 2\psi, \quad (12)$$

$$h_\times^S = h_+ \sin 2\psi - h_\times \cos 2\psi. \quad (13)$$

The end of the inspiral is handled with the exponential taper also used for the MBH-binary waveforms of Challenge 2 [4]. See directory MLDCwaveforms/FastBBH in LISAtools for the source code for these waveforms.

Data set 3.2 includes also a galactic confusion background generated from the same *detached-binary* population used in Challenge 3.1 (interacting systems have typically very small chirp masses and are not expected to make a significant contribution), but withholding all binaries with individual $A + E$ SNR > 5 , relative to instrument noise plus an estimate of confusion noise. This estimate was derived using a BIC criterion for the resolvability of individual galactic binaries [21]:

$$S_{X,\text{gal}} = 16x^2 \sin^2 x \text{ Hz}^{-1} \times \begin{cases} 10^{-44.62} (f/\text{Hz})^{-2.3} & \text{for } f \in [10^{-4}, 10^{-3}] \text{ Hz,} \\ 10^{-50.92} (f/\text{Hz})^{-4.4} & \text{for } f \in [10^{-3}, 10^{-2.7}] \text{ Hz,} \\ 10^{-62.8} (f/\text{Hz})^{-8.8} & \text{for } f \in [10^{-2.7}, 10^{-2.4}] \text{ Hz,} \\ 10^{-89.68} (f/\text{Hz})^{-20.0} & \text{for } f \in [10^{-2.4}, 10^{-2.0}] \text{ Hz} \end{cases} \quad (14)$$

(fractional frequency fluctuations, with $x = 2\pi fL$, $L \simeq 16.6782$ s). The resulting confusion background is consistent with (14), which is also used in Challenge 3 (on top of instrument noise) to define the SNRs of GW signals from MBH binaries, EMRIs (Challenge 3.3) and cosmic-string cusps (Challenge 3.4).

3.3. EMRIs

The EMRI waveforms of data set 3.3 are the Barack–Cutler [17] ‘analytic kludges’ used for Challenge 1.3.1–5 and described in [4, section 4.5], with the single change in that the number of eccentric-orbit harmonics included in the waveform does not evolve with eccentricity, but is fixed at five (lisaXML parameter `FixHarmonics`; a value of zero will reproduce the old behavior). See directory MLDCwaveforms/EMRI in LISAtools for the source code.

3.4. Cosmic string cusps

Data set 3.4 contains a number of bursts from cosmic strings, the first of two new GW sources introduced with Challenge 3. Cosmic strings are linear topological defects that may be formed

in early universe at the phase transitions predicted in many elementary-particle and superstring models. Cosmic-string oscillations emit gravitational radiation, with a substantial part of the emission from *cusps*, which can achieve very large Lorentz boosts [28]. In the limit where the tip of a cusp is moving directly toward the observer, the observed metric perturbation is a linearly polarized GW with [29]

$$h(t) = A|t - t_C|^{1/3} \times (\text{incomplete } \Gamma \text{ function envelope}), \quad A \sim \frac{G\mu L^{2/3}}{D_L}; \quad (15)$$

here t_C is the burst's central time of arrival, G is Newton's constant, μ is the string's mass per unit length, D_L is the luminosity distance to the source and L is the size of the feature that produces the cusp (e.g., the length of a cosmic string loop). If the observer's line of sight does not coincide with the cusp's direction of motion, the waveform becomes a much more complicated mixture of polarizations [30]. Under our assumption that the viewing angle departs only slightly from zero, the waveform remains dominantly linearly polarized, and the sharp spike in (15) is rounded off, introducing an exponential suppression of Fourier-domain power for frequencies above $f_{\max} = 2/(\alpha^3 L)$.

Following the model used by the LIGO Science Collaboration, we define our cusp waveforms in the Fourier domain according to

$$|h_{\times}(f)| = \mathcal{A}f^{-4/3}(1 + (f_{\text{low}}/f)^2)^{-4}, \quad h_{\times} = 0, \quad (16)$$

with $\exp(1 - f/f_{\max})$ suppression above f_{\max} . The amplitude \mathcal{A} has dimensions $\text{Hz}^{1/3}$; f_{low} sets the low-frequency cutoff of what is effectively a fourth-order Butterworth filter, which prevents dynamic-range issues with the inverse Fourier transforms (for Challenge 3 we set $f_{\text{low}} = 1 \times 10^{-5}$ Hz). The phase of the waveform is set to $\exp i(\pi - 2\pi f t_C)$ before inverse-Fourier transforming to the time domain. See directory `MLDCwaveforms/CosmicStringCusp` in `LISAtools` for the source code.

3.5. Stochastic background

Data set 3.5 contains the second GW source new to Challenge 3: an isotropic, unpolarized, Gaussian and stationary stochastic background. Allen and Romano [31] define a stochastic background as the 'gravitational radiation produced by an extremely large number of weak, independent, and unresolved gravity-wave sources, [...] stochastic in the sense that it can be characterized only statistically'. Such backgrounds are usually specified by the dimensionless quantity

$$\Omega_{\text{gw}}(f) = \frac{1}{\rho_{\text{crit}}} \frac{d\rho_{\text{gw}}}{d \log f}, \quad (17)$$

with ρ_{gw} the energy density in GWs, and $\rho_{\text{crit}} = 3c^2 H_0^2 / (8\pi G)$ the closure energy density of the universe; they are idealized as the collective, incoherent radiation of uncorrelated infinitesimal emitters distributed across the sky. If the background is isotropic, unpolarized, Gaussian and stationary, the Fourier amplitude $\tilde{h}_A(f, \hat{\Omega})$ of each emitter (with A indexing the $+$ and \times polarizations, and $\hat{\Omega}$ the direction on the 2-sphere) is completely characterized by the power-spectral-density relation [31]

$$\langle \tilde{h}_A^*(f, \hat{\Omega}) \tilde{h}_{A'}(f', \hat{\Omega}') \rangle = \frac{3H_0^2}{32\pi^3} |f|^{-3} \Omega_{\text{gw}}(|f|) \times \delta_{AA'} \delta(f - f') \delta^2(\hat{\Omega}, \hat{\Omega}'). \quad (18)$$

In Challenge 3, we assume a constant $\Omega_{\text{gw}}(f)$, as appropriate for the primordial background predicted in simple cosmological scenarios. We implement the uncorrelated emitters as a collection of 192 pseudosources distributed at *HEALPix* pixel centers across the sky. *HEALPix*

(the Hierarchical Equal Area isoLatitude Pixelization of spherical surfaces [32]) is often used to represent cosmic microwave background data sets; 192 pixels correspond to a twice-refined HEALPix grid with $N_{\text{side}} = 2^2$.

Each pseudosource consists of uncorrelated pseudorandom processes for h_+ and h_\times , generated as white noise in the time domain, and filtered to achieve the f^{-3} spectrum of (18), using the recursive $1/f^2$ filtering algorithm proposed by Plaszczyński [33], extended to spectral slope -3 . The algorithm employs a chain of $1/f^2$ infinite-impulse-response filters to reshape the white noise spectrum between minimum and maximum frequencies f_{low} and f_{knee} , set to 10^{-5} and 10^{-2} Hz in this Challenge (see the source file MLDCwaveforms/Stochastic.py in LISAtools for the Synthetic LISA implementation).

The *one-sided* PSD of each single-polarization random process (which represents the finite area of a pixel in the sky) is then given by $S_h(f)/2 = 3H_0^2/(32\pi^3)f^{-3}\Omega_{\text{gw}} \times (4\pi^2/192)$. In data set 3.5, we define $S_h^{\text{tot}} = (192 \times 2)S_h$ and we set it so that, in the TDI observables, the GW background is a few times stronger than LISA's secondary instrument noise. Namely,

$$S_h^{\text{tot}}(f) = 0.7\text{--}1.3 \times 10^{-47} (f/\text{Hz})^{-3} \text{ Hz}^{-1} \quad (19)$$

(taking $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$, this corresponds to $\Omega_{\text{gw}} = 8.95 \times 10^{-12}\text{--}1.66 \times 10^{-11}$).

One of the more promising approaches to detect GW backgrounds with LISA relies on estimating instrument noise levels by way of symmetrized TDI observables that are insensitive to GWs at low frequencies in the LISA band [34–36]. For realistic LISA orbits, however, the low-frequency behavior of such observables becomes more complicated than discussed in the literature. To simplify the initial investigation of the background-detection problem in data set 3.5, we have therefore approximated LISA as a rigidly rotating triangle with equal and constant armlengths (i.e., Synthetic LISA's CircularRotating).

4. Conclusion

Since their inception, the Mock LISA Data Challenges have received remarkable support from the GW community, and have set the stage for many practical demonstrations of the feasibility of LISA's exciting science with present-day data-analysis techniques. Future challenges will feature ever more realistic models of waveforms and instrument noise, and they will endeavour to scope out all important aspects of the LISA science objectives. In addition, the software tools developed for the MLDCs [16] can be used to generate data sets for many other data-analysis experiments outside the main challenges; and the MLDC *standard model* of LISA's observations (including the MLDC 'pseudo-LISA' and GW models) is proving extremely valuable to the current analytical investigations of the LISA science performance that are being run by the LIST.

To obtain more information and to participate in the MLDCs, see the official MLDC website (astrogravs.nasa.gov/docs/mldc) and the Task Force wiki (www.tapir.caltech.edu/listwg1b).

Acknowledgments

SB and EP were supported by the Deutsches Zentrum für Luft- und Raumfahrt; LW's work by the Alexander von Humboldt Foundation's Sofja Kovalevskaja Programme funded by the German Federal Ministry of Education and Research. JG thanks the Royal Society for support and the Albert Einstein Institute for hospitality and support while part of this work was being completed. SF was supported by the Royal Society, BSS and IH by the UK Science and Technology Facilities Council. MV was supported by the LISA Mission Science Office and

by JPL's HRDF. CC's, JC's and MV's work was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration. MT acknowledges support from the Spanish Ministerio de Educación y Ciencia Research projects FPA-2007-60220, CSD207-00042 and the Govern de les Illes Balears, Conselleria d'Economia, Hisenda i Innovació. IM was partially supported by NASA ATP grant NNX07AH22G to Northwestern University.

References

- [1] Bender P and Danzmann P (the LISA Study Team) 1998 *Laser Interferometer Space Antenna for the Detection of Gravitational Waves, Pre-Phase A Report* **MPQ 233** (Garching: Max-Planck-Institut für Quantenoptik)
- [2] Arnaud K A *et al* (the MLDC Task Force) 2006 *Laser Interferometer Space Antenna: 6th Int. LISA Symp. (Greenbelt, MD, 19–23 Jun 2006)* ed S M Merkowitz and J C Livas (Melville, NY: AIP) p 619
- Arnaud K A *et al* (the MLDC Task Force) 2006 *Laser Interferometer Space Antenna: 6th International LISA Symp. (Greenbelt, MD, 19–23 Jun 2006)* ed S M Merkowitz and J C Livas (Melville, NY: AIP) p 625 (long version with lisaXML description: *Preprint gr-qc/0609106*)
- [3] Arnaud K A *et al* (the MLDC Task Force and Challenge 1 participants) 2007 *Class. Quantum Grav.* **24** S529
- [4] Arnaud K A *et al* (the MLDC Task Force) 2007 *Class. Quantum Grav.* **24** S551
- [5] Babak S *et al* (the MLDC Task Force and Challenge 2 Participants) 2008 *Class. Quantum Grav.* **25** 114037
- [6] Vallisneri M, Crowder J and Tinto M 2008 *Class. Quantum Grav.* **25** 065005
- [7] Xspec, an X-ray Spectral Fitting Package, heasarc.gsfc.nasa.gov/docs/xanadu/xspec
- [8] Jaranowski P, Krolak A and Schutz B F 1998 *Phys. Rev. D* **58** 063001
- [9] Whelan J T, Prix R and Khurana D 2008 Improved search for galactic white dwarf binaries in Mock LISA Data Challenge 1B using an F-statistic template bank *Class. Quantum Grav.* **25** 184029 (*Preprint arXiv:0805.1972*)
- [10] Trias M, Vecchio M and Veitch J 2008 Markov chain Monte Carlo searches for galactic binaries in MLDC-1B data sets *Class. Quantum Grav.* **25** 184028 (*Preprint arXiv:0804.4029*)
- [11] Gijs Nelemans' LISA wiki, www.astro.kun.nl/nelemans/dokuwiki
- [12] Harry I W, Fairhurst S and Sathyaprakash B S 2008 A hierarchical search for gravitational waves from supermassive black hole binary mergers *Class. Quantum Grav.* **25** 184027 (*Preprint arXiv:0804.3274*)
- [13] Gair J R, Mandel I and Wen L 2008 Improved time–frequency analysis of extreme-mass-ratio inspiral signals in Mock LISA data *Class. Quantum Grav.* **25** 184031 (*Preprint arXiv:0804.1084*)
- [14] Gair J R, Babak S, Porter E K and Barack L 2008 A constrained Metropolis–Hastings search for EMRIs in the Mock LISA Data Challenge 1B *Class. Quantum Grav.* **25** 184030 (*Preprint arXiv:0804.3322*)
- [15] Cornish N J 2008 Detection strategies for extreme mass ratio inspirals *Preprint arXiv:0804.3323*
- [16] LISAtools website and SVN repository, lisatools.googlecode.com
- [17] Barack L and Cutler C 2004 *Phys. Rev. D* **69** 082005
- [18] Cornish N J and Rubbo L J 2003 *Phys. Rev. D* **67** 022001
- Cornish N J and Rubbo L J 2003 *Phys. Rev. D* **67** 029905 (erratum)
- LISA Simulator website, www.physics.montana.edu/lisa; now included in [16]
- [19] Vallisneri M 2005 *Phys. Rev. D* **71**
- Synthetic LISA website, www.vallis.org/syntheticlisa; now included in [16]
- [20] Petiteau A, Auger G, Halloin H, Jeannin O, Plagnol E, Pireaux S, Regimbeau T and Vinet J-Y 2008 *Phys. Rev. D* **77** 023002
- LISACode website, www.apc.univ-paris7.fr/LISA-France/analyse.phtml; now included in [16]
- [21] Cornish N J and Littenberg T B 2007 *Phys. Rev. D* **76** 083006
- [22] Nelemans G, Yungelson L R and Portegies Zwart S F 2001 *Astron. Astrophys.* **375** 890
- [23] Nelemans G, Yungelson L R and Portegies Zwart S F 2004 *Mon. Not. R. Astron. Soc.* **349** 181
- [24] Roelofs G H A, Nelemans G and Groot P J 2007 *Mon. Not. R. Astron. Soc.* **382** 685
- [25] Damour T, Iyer B and Sathyaprakash B S 2001 *Phys. Rev. D* **63** 044023
- [26] Apostolatos T A, Cutler C, Sussman G J and Thorne K S 1994 *Phys. Rev. D* **49** 6274
- [27] Lang R N and Hughes S A 2006 *Phys. Rev. D* **74** 122001
- [28] Damour T and Vilenkin A 2000 *Phys. Rev. Lett.* **85** 3761
- [29] Siemens X, Creighton J, Maor I, Ray Majumder S, Cannon K and Read J 2006 *Phys. Rev. D* **73** 105001
- [30] Siemens X and Olum K D 2003 *Phys. Rev. D* **68** 085017
- [31] Allen B and Romano J D 1999 *Phys. Rev. D* **59** 102001

- [32] Górski K M *et al* 2005 *Astrophys. J.* **622** 759
HEALPix website, healpix.jpl.nasa.gov
- [33] Plaszczyński S 2007 *Fluc. Noise Lett.* **7** R1
- [34] Tinto M, Armstrong J W and Estabrook F B 2000 *Phys. Rev. D* **63** 021101
- [35] Hogan C J and Bender P L 2001 *Phys. Rev. D* **64** 062002
- [36] Robinson E L, Romano J D and Vecchio A 2008 Search for a stochastic gravitational-wave signal in the second round of the Mock LISA Data Challenges *Class. Quantum Grav.* **25** 184019 (Preprint [arXiv:0804.4144](https://arxiv.org/abs/0804.4144))